

LncRNA NNT-AS1 Affects Progesterone Resistance by Regulating MiR-542-3p/survivin Axis in Endometrial Cancer

Chen Guo^{a,*}, Wei Zhao^a, Xiu-fang Bao^a

^aAffiliated Hospital of Inner Mongolia University Nationalities. Department Obstetrics & Gynecology. Tong Liao 028000.

Abstract

Background: Progestin is commonly used for young patients suffering from endometrial cancer, but more than 30% of patients with progestin treatment have presented progestin resistance. This study aimed to investigate the involvement of NNT-AS1 in progesterone resistance endometrial cancer and assess its possible molecular mechanism.

Methods: Establishment of progesterone resistant Ishikawa (Ishikawa-PR) cell line that are resistant to the growth-inhibitory effects of progestin in vitro. Protein and mRNA expression were determined by western blot and qRT-PCR, respectively. MTT assay tested cell proliferation. Silence and overexpression of NNT-AS1 and survivin were performed by si-NNT-AS1/survivin and pcDNA-NNT-AS1/survivin, respectively. The effect of NNT-AS1 on Ishikawa-PR cell transplant tumor growth was measured by mouse xenograft model. In addition, we evaluated the relationship between NNT-AS1 and miR-542-3p by RNA immunoprecipitation (RIP) and RNA pull down.

Results: The NNT-AS1 and survivin expression were both significantly increased, while miR-542-3p was decreased in Ishikawa-PR cell. Overexpression of NNT-AS1 was found to increase sensitive Ishikawa cell resistance, while inhibition of NNT-AS1 could reduce Ishikawa-PR cell resistance. NNT-AS1 functioned as a miR-542-3p decoy, and miR-542-3p could regulate Ishikawa cell resistance by survivin. In addition, NNT-AS1 was confirmed to regulate survivin expression and Ishikawa cell resistance by miR-542-3p. In vivo mice xenograft model revealed that silencing NNT-AS1 could improve endometrial cancer resistance.

Conclusion: LncRNA NNT-AS1 affect progesterone resistance by regulating miR-542-3p/survivin axis in endometrial cancer.

Keywords: LncRNA NNT-AS1; progesterone resistance; endometrial cancer; miR-542-3p; survivin

INTRODUCTION

Endometrial cancer is the most common gynecological tumor in developed countries. Hormonal therapy, such as medroxyprogesterone acetate (MPA), has been applied in the conservative treatment of young patients who wish to preserve their fertility, as well as in the palliative treatment of advanced-stage patients [1]. But more than 30% of patients with progestin treatment have presented progestin resistance [2]. In recent years, there is some improvement has been demonstrated in endometrial cancer progesterone resistance. However, the mechanisms underlying acquired resistance to progestin is not entirely clear.

Long noncoding RNAs (lncRNAs) are over 200 nucleotides in length without protein-coding capacity. Guo et al [3] demonstrated that GAS5 acted as a tumor suppressor lncRNA in endometrial cancer, and could enhance PTEN expression to promote cancer cell apoptosis by inhibiting miR-103. Huang et al [4] reported that HOTAIR expression was increased in endometrial cancer cells and tissues, and the down-regulation of HOTAIR resulted in a significant inhibition of cell proliferation, migration, and invasion and in cell cycle arrest at the G0/G1 phase. Furthermore, HOTAIR depletion significantly suppressed the endometrial cancer tumorigenesis in vivo. On top of this, there are several lncRNA that play an important role in endometrial carcinoma development, such as ASLNC04080 [5], FER1L4 [6], BANC1 [7], MEG3 [8], etc. LncRNA Nicotinamide Nucleotide Transhydrogenase-antisense RNA1 (NNT-AS1), which located in 5p12 with 3 exons, has been mapped to chromosome 5 region 43573185–43603230 according to the NCBI (GRCh38.p2). Two recent articles by Wang [9] and Hua [10] have respectively reported that NNT-AS1 was increased in col-

*Corresponding author: Chen Guo
Mailing address: Affiliated Hospital of Inner Mongolia University Nationalities. Department Obstetrics & Gynecology. Tong Liao 028000, China.
E-mail: koala7@126.com
Received: 29 March 2019 Accepted: 10 June 2019

orectal cancer and cervical cancer, and associated with advanced clinicopathological features and poor overall survival. However, NNT-AS1 have not been investigated in endometrial cancer, especially in progesterone resistance endometrial cancer.

Survivin, the smallest member of IAP (inhibitor of apoptosis) family, is a dual functional protein acting as a critical apoptosis inhibitor and key cell cycle regulator. Survivin is reported to highly express in most human cancers, such as lung, pancreatic and breast cancers [11]. Also, survivin plays an essential role in the drug resistant phenotype of multiple human cancers [12,13]. Especially, survivin was involved in progesterone resistance endometrial cancer/hyperplasia [14,15]. Overexpression of survivin was found in progestin-resistant endometrial samples, and down regulation of survivin increased the progestin sensitivity in endometrial cancer. A number of studies have shown that many miRNAs regulated cancer cell function and drug resistance by targeting survivin, for example miR-218 [16], miR-214-3p [17], miR-138-5p [18], etc. Althoff et al [19] using reporter assays confirmed

miR-542-3p directly targeted survivin in neuroblastoma. In addition, miR-542-3p played an important role in endometrial decidualization by regulating IGFBP1 [20]. Therefore, we hypothesized that miR-542-3p may regulate endometrial cancer cell resistance by targeting survivin.

Based on the result of predictive analytics software, NNT-AS1 may function as a miR-542-3p decoy. Thus, this study aimed to investigate the expression of NNT-AS1 and miR-542-3p and evaluate whether NNT-AS1 affect progesterone resistance by regulating miR-542-3p/survivin axis in progesterone resistance endometrial cancer.

MATERIALS AND METHODS

Routine cell culture and establishment of progestin-resistant cell line

Ishikawa cells were grown in DMEM Medium (Invitrogen) containing 10% FBS (Gibco), 50 mg/mL gentamycin. Progestin-resistant Ishikawa cell lines were induced as previous studies [21]. The progestin-resistant Ishikawa cell were grown in DMEM Medium (Invitrogen) contain-




Figure 1. The increased NNT-AS1 expression in progesterone-resistant Ishikawa cells. **(A)** The growth of the two cells was tested by MTT assay and analyzed by plotting a cell growth curve. **(B)** The NNT-AS1 expression was significantly increased in Ishikawa-PR. **(C)** The miR-542-3p expression was significantly decreased in Ishikawa-PR. **(D)** The survivin expression was increased in Ishikawa-PR. * $P < 0.05$ vs Ishikawa.




Figure 2. The effect of NNT-AS1 on Ishikawa cell resistance. **(A)** Overexpression NNT-AS1 promoted cell resistance in sensitive Ishikawa cells. **(B)** Knockdown of NNT-AS1 reduced cell resistance in progesterone-resistant Ishikawa cells. *P<0.05 vs si-control or pcDNA.

ing 10% FBS (Gibco), 50 mg/mL gentamycin and maintained with a kind of synthetic progestin medoxyprogesterone acetate (MPA, 10 μM, Sigma).

MTT assay

Seven-day growth curves of the parent Ishikawa cells and the progestin-resistant Ishikawa cells were tested by MTT assay. Cells were plated in a 96 well flat bottomed microplate with 100 μl cell suspension (1~2*10⁴ cells/ml) per well. The parent Ishikawa cells were grown in the complete medium with absence of MPA. The progestin-resistant Ishikawa cells were routinely maintained in the complete medium containing 10 μM MPA. Culture media were changed every other day.

After transfection, Ishikawa cell was stimulated with 1 μM MPA, while progestin-resistant Ishikawa cell were stimulated with 5 μM MPA. And then cell proliferation was detected by MTT assay. A total of 10 μl MTT (5 mg/ml) was added into each well. After incubation for 4h, the medium was poured off, and formazan crystals were dissolved with 150 μl DMSO by shaking. The absorbance was measured at 490 nm with a microplate reader.

Cell transfection

MiR-542-3p mimic/inhibitor or negative control mimic/inhibitor were purchased from Genepharma, and the NNT-AS1 and survivin expression plasmids were also constructed by Genepharma (Shanghai, China). Si-NNT-AS1, si-survivin and their respective negative control were chemically synthesized by RiboBio (Shanghai, China). Ishikawa cells (with progestin-resistant) were transfected with NNT-AS1 and survivin siRNA or expression plasmids using Lipofectamine 2000 (Invitrogen). To overexpress or knockdown miR-542-3p in Ishikawa cells (progestin-resistant), the cells were transfected with miR-542-3p mimics or inhibitor using

Lipofectamine 2000 (Invitrogen). After transfection, Ishikawa cell was stimulated with 1 μM MPA, while progestin-resistant Ishikawa cell were stimulated with 5 μM MPA.

Western blot

The total protein was extracted from tissues and cells. Equal amounts of proteins were fractioned with SDS-PAGE followed by electrophoretic transfer of proteins onto nitrocellulose membranes. The blots were probed with antibodies against MDR-1 and survivin (Abcam), and followed by incubation with secondary antibodies conjugated with horseradish peroxidase (ThermoFisher). The immune complexes were detected with a WesternBright™ ECL Western Blotting HRP Substrate kit and analyzed with image lab software (BioRad, USA).

RNA extraction and qRT-PCR analysis

Total RNA from tissues and cells were extracted using Trizol reagent (TAKARA). RNA was reversely transcribed into cDNAs using the Primer-Script one step RT-PCR kit (TAKARA, Dalian, China). The cDNA template was amplified by real-time RT-PCR using the SYBR Premix Dimmer Eraser kit (TAKARA). Gene expression in each sample was normalized to β-actin expression. Real-time PCR reactions were performed using the ABI7500 system (Applied Biosystems, USA). The real-time PCRs were performed in triplicate. Relative expression fold change of mRNAs was calculated by the 2^{-ΔΔCt} method.

RNA pull-down

To determine whether NNT-AS1 is associated with the RISC complex, we performed an RNA precipitation assay by synthesized NNT-AS1 as a probe and then detected AGO2 from the pellet by western blot and miR-542-3p by qRT-PCR. The biotin-labeled full length lncRNA NNT-AS1




Figure 4. The miR-542-3p regulated Ishikawa cell resistance by survivin. **(A)** miR-542-3p inhibitor increased sensitive Ishikawa cell resistance and upregulated MDR-1 expression, while si-survivin reversed the effect of the miR-542-3p inhibitor. **(B)** miR-542-3p mimic reduced Ishikawa-PR cell resistance and MDR-1 expression, pcDNA-survivin reversed the effect of miR-542-3p mimic. *P<0.05 vs NC or pre-NC, #P <0.05 vs miR-542-3p inhibitor +si-control or miR-542-3p mimic + pcDNA.

BALB/c mice used in the present study were treated according to the protocols approved by the ethical committee of the Affiliated Hospital of Inner Mongolia University Nationalities. Mice at the age of 4–6 weeks (n =6) were injected subcutaneously with siRNA-NNT-AS1 transfected- Ishikawa-PR cells (5×10^6 in the 100 ul PBS) in the right and left hind limbs. The control mice (n =6) were injected with the same volume of vector only. Seven days after injection, the mice received MPA (100 mg/kg body weight), which was injected three times/week (nine times in total). The mice were killed on day 28 after tumor cell transplantation. Tumor size was measured with calipers weekly, and tumor volume was calculated using the following formula: tumor volume (mm³) = (tumor length * tumor width ²)/2. All experimental procedures were in accordance with the National Institute of Health Guide for the Care and Use of Laboratory Animals.

Statistical analysis

Data analysis was performed using SPSS 17.0. Data were represented as mean ± SD based on at least three repeats. Group difference was assessed using Student's t test, and P<0.05 was considered as statistically significant.

RESULTS

The increased lncRNA NNT-AS1 expression in progesterone-resistant Ishikawa cells

To establish progestin resistant endometrial cell line (Ishikawa-PR), we used MPA, a kind of synthetic progestin medoxyprogesterone acetate to induce Ishikawa cell as described in the literature [22]. Cells were plated in 96-well flat-bottomed microplates as described in Materials and methods. The 7 day growth curve of Ishikawa cells and the progestin-resistant Ishikawa cells shown




Figure 5. NNT-AS1 regulated survivin expression by miR-542-3p. **(A)** pcDNA-NNT-AS1 increased survivin mRNA and protein expression, while miR-542-3p mimic reversed the effect of pcDNA-NNT-AS1 in sensitive Ishikawa cell. **(B)** siRNA-NNT-AS1 suppressed survivin mRNA and protein expression, while the miR-542-3p inhibitor reversed the effect of si-NNT-AS1 in Ishikawa-PR cell. * $P < 0.05$ vs pcDNA or si-control, # $P < 0.05$ vs pcDNA-NNT-AS1+pre-NC or siRNA-NNT-AS1+NC.

in Figure 1A, which suggested a successfully established experimental progesterone-resistant Ishikawa cell. And then the expression levels of lncRNA NNT-AS1, miR-542-3p and survivin in the two cells were analyzed by RT-qPCR or western blot. The RT-qPCR results showed that lncRNA NNT-AS1 expression was significantly increased, while the miR-542-3p expression was obviously decreased in Ishikawa-PR cells (Figure 1B and C). The western blot results showed that survivin expression was also significantly increased in Ishikawa-PR cells (Figure 1D). These data revealed that the expression of lncRNA NNT-AS1, miR-542-3p and survivin were differ-

ent in sensitive Ishikawa cell line and Ishikawa-PR cells, and the lncRNA NNT-AS1 and survivin expression were significantly increased while miR-542-3p was decreased in progesterone-resistant Ishikawa cells.

The effect of lncRNA NNT-AS1 on Ishikawa cell resistance

To investigate the effect of overexpressing exogenous and RNA-interfered endogenous lncRNA NNT-AS1 on Ishikawa cell resistance, the sensitive Ishikawa cell transfection of pcDNA-NNT-AS1 was stimulated by MAP (0, 0.01, 0.1, 1, 2 μ M), and the Ishikawa-PR cells transfection of si-NNT-AS1 was stimulated by MAP (0,




Figure 6. NNT-AS1 regulated Ishikawa cell resistance by miR-542-3p. **(A)** pcDNA-NNT-AS1 increased cell resistance and MDR-1 expression, while the miR-542-3p mimic reversed the effect of pcDNA-NNT-AS1 in sensitive Ishikawa cell. **(B)** siRNA-NNT-AS1 reduced cell resistance and MDR-1 expression, while miR-542-3p inhibitor reversed the effect of siRNA-NNT-AS1 in Ishikawa-PR cell. *P<0.05 vs pcDNA or si-control, #P <0.05 vs pcDNA-NNT-AS1+pre-NC or siRNA-NNT-AS1+NC.

0.5, 1.0, 5.0, 10 μ M). The MTT assay showed that when the MAP concentration was greater than 0.1 μ M, the cell proliferation in pcDNA-NNT-AS1-transfected cell was significantly higher than that in pcDNA-transfected cell, suggesting overexpression NNT-AS1 promoted cell resistance in sensitive Ishikawa cells (Figure 2A). Moreover, when the MAP concentration was greater than 5 μ M, the cell proliferation in si-NNT-AS1-transfected cell was significantly lower than in si-control-transfected cell, suggesting knockdown of NNT-AS1 reduced cell resist-

ance in Ishikawa-PR cells (Figure 2B). These results indicated that lncRNA NNT-AS1 played an important role in Ishikawa cell resistance.

LncRNA NNT-AS1 functioned as a miR-542-3p decoy in Ishikawa cells

As the NNT-AS1 expression was opposite to miR-542-3p expression, we suspected that NNT-AS1 may function as a miR-542-3p decoy. The Figure 3A displayed the predicted positions of miR-542-3p binding sites on




Figure 7. RNA interference of NNT-AS1 improved endometrial cancer resistance in mice. **(A)** siRNA-NNT-AS1 minimized endometrial cancer resistance in mice. In siRNA-NNT-AS1 transfected-Ishikawa-PR cell transplantation mice: **(B)** The NNT-AS1 expression was significantly decreased. **(C)** The miR-542-3p expression was significantly increased. **(D)** The MDR-1 and survivin expression were both decreased. * $P < 0.05$ vs si-control.

NNT-AS1 transcript, and then their relationship was confirmed by RNA immunoprecipitation and RNA pull-down assay. The AGO2 antibody was used into RNA immunoprecipitation, and the RT-PCR results showed that, compared to IgG complex, the NNT-AS1 and miR-542-3p were both existed and obviously increased in AGO2 complex (Figure 3B). The RNA pull-down result showed that the AGO2 existed in the NNT-AS1 drop-down compound, which confirmed a direct interaction of NNT-AS1 with miR-542-3p (Figure 3C). The miR-542-3p expression was significantly increased in the NNT-AS1 drop-down compound, which further confirmed NNT-AS1 functioned as a miR-542-3p decoy (Figure 3D). These data suggested that NNT-AS1 could function as a miR-542-3p decoy in Ishikawa cells.

The miR-542-3p could regulate Ishikawa cell resistance by survivin

A previous study has reported that miR-542-3p exert-

ed tumor suppressive functions by targeting survivin in neuroblastoma^[22], and survivin expression was opposite to miR-542-3p expression in our study. So, we speculated that miR-542-3p could regulate Ishikawa cell resistance by targeting survivin. In sensitive Ishikawa cells, transfection of miR-542-3p inhibitor increased cell resistance and MDR-1 expression, however, co-transfection with si-survivin reversed the increased cell resistance and MDR-1 expression (Figure 4A). In Ishikawa-PR cells, transfection of miR-542-3p mimic reduced cell resistance and MDR-1 expression, however, co-transfection with pcDNA-survivin reversed the reduced cell resistance and MDR-1 expression (Figure 4B). These data implied that miR-542-3p could regulate Ishikawa cell resistance by regulating survivin.

LncRNA NNT-AS1 regulated survivin expression by miR-542-3p

To explore whether the survivin expression could be reg-

ulated by NNT-AS1 through miR-542-3p, the two cells were cotransfected with overexpressing exogenous and RNA-interfered endogenous NNT-AS1 and miR-542-3p mimic and inhibitor, respectively. As shown in Figure 5A, sensitive Ishikawa cells transfected with pcDNA-NNT-AS1 upregulated survivin mRNA and protein expression, while co-transfection with miR-542-3p mimic reversed the increased survivin expression. As Figure 5B showed that Ishikawa-PR cells transfected with siRNA-NNT-AS1 downregulated survivin mRNA and protein expression, while co-transfection with miR-542-3p inhibitor reversed the decreased survivin expression. These results suggested that lncRNA NNT-AS1 could regulate survivin expression by miR-542-3p in Ishikawa cells.

LncRNA NNT-AS1 regulated Ishikawa cell resistance by miR-542-3p

Next, to demonstrate miR-542-3p was involved in the regulation mechanism of Ishikawa cell resistance by lncRNA NNT-AS1, the two cells were cotransfected with overexpressing exogenous and RNA-interfered endogenous NNT-AS1 and miR-542-3p mimic and inhibitor, respectively. As shown in Figure 6A, sensitive Ishikawa cells transfected with pcDNA-NNT-AS1 increased cell resistance and upregulated MDR-1 expression, while co-transfection with miR-542-3p mimic reversed the effect of pcDNA-NNT-AS1. As Figure 6B showed that Ishikawa-PR cells transfected with siRNA-NNT-AS1 reduced cell resistance and downregulated MDR-1 expression, while co-transfection with miR-542-3p inhibitor reversed the effect of siRNA-NNT-AS1. These results suggested that lncRNA NNT-AS1 could regulate Ishikawa cell resistance by miR-542-3p.

Knockdown of NNT-AS1 improved endometrial cancer resistance in mice

To further study the role of RNA-interfered endogenous NNT-AS1 in the improvement of endometrial cancer resistance in vivo, siRNA-NNT-AS1 transfected-Ishikawa-PR cells were injected subcutaneously into the right and left hind limbs of mice, and 7 days after injection, the mice received MPA. According to the measured tumor size, the siRNA-NNT-AS1 group had a smaller tumor than that in si-control group, which indicated that siRNA-NNT-AS1 minimized endometrial cancer resistance in mice (Figure 7A). As Figure 7B showed that, the NNT-AS1 expression in tumor tissues that isolated from siRNA-NNT-AS1 transfected-progestin resistant Ishikawa cells transplantation mice was significantly lower than that from si-control transfected-progestin resistant Ishikawa cells transplantation mice. The miR-542-3p expression was obviously increased in siRNA-NNT-AS1 transfected-progestin resistance Ishikawa cells transplantation mice (Figure 7C). In addition, the survivin and

MDR-1 protein expression were both decreased in siRNA-NNT-AS1 transfected-progestin resistance Ishikawa cells transplantation mice, which further suggested that siRNA-NNT-AS1 could minimize endometrial cancer resistance in mice (Figure 7D). These results demonstrated that RNA-interfered endogenous NNT-AS1 could improve endometrial cancer resistance in mice.

DISCUSSION

Progestin resistance is the main obstacle to successful conservative therapy in young endometrial cancer patients. To investigate the molecular events that lead to progestin resistance and to find a possible way to reverse progestin resistance in endometrial cancer, we established an Ishikawa-PR cell line that was resistant to the growth-inhibitory effects of progestin in vitro. The 7 day growth curve of Ishikawa cells and the Ishikawa-PR cell (Figure 1A) indicated a successfully established experimental progesterone-resistant Ishikawa cell line. Survivin was involved in progesterone resistance endometrial cancer/hyperplasia^[14]. In the present study, we found survivin protein expression was increased in Ishikawa-PR (Figure 1D), and inhibition of survivin increased the progestin sensitivity and decreased MDR-1 expression (Figure 4A), which was in agreement with previously reported results^[15].

MiRNAs are small RNA molecules (22 nt) that interact with their target mRNAs inhibiting translation or/and cleaving the target mRNA. Studies by Yoon^[23] and Althoff^[19] both demonstrated that miR-542-3p could directly target survivin in A549 cells and neuroblastoma cell lines, respectively. Meanwhile, miR-542-3p was increased in pairs ectopic/eutopic endometrium as compared with normal endometrium, and played an important role in endometrial decidualization^[20]. In our study, miR-542-3p was significantly reduced in Ishikawa-PR (Figure 1C). The miR-542-3p inhibitor was found to increase sensitive Ishikawa cell resistance and MDR-1 expression, and miR-542-3p mimic reduced Ishikawa-PR cell resistance and MDR-1 expression. Moreover, in the Ishikawa cell, si-survivin reversed the miR-542-3p inhibitor-increased cell resistance and MDR-1 expression, and in Ishikawa-PR cell, pcDNA-survivin reversed the miR-542-3p mimic-reduced cell resistance and MDR-1 expression (Figure 4). The data presented here suggested that miR-542-3p could regulate Ishikawa cell resistance by survivin.

Increasing lines of evidence show that lncRNAs are important factors in cancer and confirmed that lncRNAs might function as a competing endogenous RNA or a molecular sponge in modulating miRNA^[24]. Li et al^[25] using luciferase reporter and qRT-PCR assays confirmed that miR-200c bound directly to MALAT1, and disrupting MALAT1/miR-200c sponge decreased invasion and

migration in endometrioid endometrial carcinoma. This study, using RIP and RNA pull-down assay provided the first evidence that NNT-AS1 functioned as a miR-542-3p decoy in Ishikawa cells (Figure 3). Furthermore, overexpression NNT-AS1 was found in colorectal cancer and cervical cancer by literature, and NNT-AS1 inhibition could suppress cancer cells proliferation and invasion ability in vitro^[9,10]. In this study, NNT-AS1 was obviously higher in Ishikawa-PR than in sensitive Ishikawa cell (Figure 1B). Further studies have revealed that overexpression NNT-AS1 promoted cell resistance in sensitive Ishikawa cells, but the knockdown of NNT-AS1 reduced cell resistance in Ishikawa-PR (Figure 2).

In addition, we found that pcDNA-NNT-AS1 upregulated survivin expression which could be reversed by miR-542-3p, and siRNA-NNT-AS1 downregulated survivin expression which could be reversed by miR-542-3p inhibitor, suggesting that NNT-AS1 regulated survivin expression by miR-542-3p (Figure 5). Further, we also found that pcDNA-NNT-AS1 increased cell resistance and upregulated MDR-1 expression which could be reversed by miR-542-3p in sensitive Ishikawa cells, and siRNA-NNT-AS1 reduced cell resistance and downregulated MDR-1 expression which could be reversed by miR-542-3p inhibitor in the Ishikawa-PR cell, suggesting that NNT-AS1 regulated survivin expression by miR-542-3p (Figure 6). In consideration of the above-described biological behaviors of NNT-AS1, to further investigate the role of NNT-AS1 in the improvement of endometrial cancer resistance in vivo, we utilized siRNA-mediated knockdown of NNT-AS1 and evaluated that the resultant effects on tumor growth. We found that silencing NNT-AS1 obviously improved the sensitivity of Ishikawa-PR cell transplant tumors to MAP, and the expression level changes of miR-542-3p, survivin and MDR-1 were in accordance with in vitro (Figure 7). All these data implied that NNT-AS1 affected progesterone resistance by the miR-542-3p/survivin axis in vitro and in vivo.

In conclusion, our data showed an abnormal expression of lncRNA NNT-AS1 in progesterone resistance Ishikawa cell, and demonstrated that the inhibition of NNT-AS1 reduced progesterone resistance by regulating miR-542-3p/survivin axis. Therefore, NNT-AS1 is a potential therapeutic target for the treatment of progesterone resistance endometrial cancer.

COMPETING INTERESTS

The authors have declared that they have no competing interests.

REFERENCES

1. Park, J. Y., and Nam, J. H. (2015) Progesterins in the Fertility-Sparing Treatment and Retreatment of Patients With Primary and Recurrent Endometrial Cancer. *Oncologist* 20, 270-278
2. Gunderson, C. C., Fader, A. N., Carson, K. A., and Bristow, R. E. (2012) Oncologic and Reproductive outcomes with progestin therapy in women with endometrial hyperplasia and grade 1 Adenocarcinoma: A systematic review. *Gynecologic Oncology* 125, 477-482
3. Guo, C., Song, W. Q., Sun, P., Jin, L., and Dai, H. Y. (2015) LncRNA-GAS5 induces PTEN expression through inhibiting miR-103 in endometrial cancer cells. *Journal of Biomedical Science* 22, 100
4. Huang, J., Ke, P., Guo, L., Wang, W., Tan, H., Liang, Y., and Yao, S. (2014) Lentivirus-mediated RNA interference targeting the long noncoding RNA HOTAIR inhibits proliferation and invasion of endometrial carcinoma cells in vitro and in vivo. *International Journal of Gynecological Cancer* 24, 635-642
5. Zhai, W., Li, X., Wu, S., Zhang, Y., Pang, H., and Chen, W. (2015) Microarray expression profile of lncRNAs and the upregulated ASLNC04080 lncRNA in human endometrial carcinoma. *International Journal of Oncology* 46, 2125-2137
6. Qiao, Q., and Hong, L. (2016) LncRNA FER1L4 suppresses cancer cell proliferation and cycle by regulating PTEN expression in endometrial carcinoma. *Biochemical & Biophysical Research Communications* 478, 507-512
7. Wang, D., Wang, D., Wang, N., Long, Z., and Ren, X. (2016) Long Non-Coding RNA BANCR Promotes Endometrial Cancer Cell Proliferation and Invasion by Regulating MMP2 and MMP1 via ERK/MAPK Signaling Pathway. *Cellular Physiology & Biochemistry* 40, 644
8. Guo, Q., Qian, Z., Yan, D., Li, L., and Huang, L. (2016) LncRNA-MEG3 inhibits cell proliferation of endometrial carcinoma by repressing Notch signaling. *Biomedicine & Pharmacotherapy* 82, 589-594
9. Qian, W., Lei, Y., Xin, H., Jiang, Y., Hu, Y., Zhe, L., Jian, L., Tao, W., Ma, Y., and An, G. (2017) Upregulated NNT-AS1, a long noncoding RNA, contributes to proliferation and migration of colorectal cancer cells in vitro and in vivo. *Oncotarget* 8, 3441-3453
10. Hua, F., Liu, S., Zhu, L., Ma, N., Jiang, S., and Yang, J. (2017) Highly expressed long non-coding RNA NNT-AS1 promotes cell proliferation and invasion through Wnt/ β -catenin signaling pathway in cervical cancer. *Biomedicine & Pharmacotherapy* 92, 1128-1134
11. Chen, X., Duan, N., Zhang, C., and Zhang, W. (2016) Survivin and Tumorigenesis: Molecular Mechanisms and Therapeutic Strategies. *Journal of Cancer* 7, 314-323
12. Kar, R., Palanichamy, J. K., Banerjee, A., Chattopadhyay, P., Jain, S. K., and Singh, N. (2015) Survivin siRNA increases sensitivity of primary cultures of ovarian cancer cells to paclitaxel. *Clinical & Translational Oncology* 17, 737-742
13. Berinstein, N. L., Karkada, M., Oza, A. M., Odunsi, K., Vilella, J. A., Nemunaitis, J. J., Morse, M. A., Pejovic, T., Bentley, J., and Buyse, M. (2015) Survivin-targeted immunotherapy

- drives robust polyfunctional T cell generation and differentiation in advanced ovarian cancer patients. *Oncoimmunology* 4, e1026529
14. Chen, X., Zhang, Z., Feng, Y., Fadare, O., Wang, J., Ai, Z., Jin, H., Gu, C., and Zheng, W. (2009) Aberrant survivin expression in endometrial hyperplasia: another mechanism of progesterin resistance. *Modern Pathology An Official Journal of the United States & Canadian Academy of Pathology Inc* 22, 699-708
 15. Fan, R., Wang, Y., Yue, W., Li, W., and Zheng, W. (2017) Mechanism of progesterin resistance in endometrial precancer/cancer through Nrf2-survivin pathway. *American Journal of Translational Research* 9, 1483-1491
 16. Hu, Y., and Xu, K. (2015) miR-218 targets survivin and regulates resistance to chemotherapeutics in breast cancer. *Breast Cancer Research & Treatment* 151, 269-280
 17. Phatak, P., Byrnes, K. A., Mansour, D., Liu, L., Cao, S., Li, R., Rao, J. N., Turner, D. J., Wang, J. Y., and Donahue, J. M. (2016) Overexpression of miR-214-3p in esophageal squamous cancer cells enhances sensitivity to cisplatin by targeting survivin directly and indirectly through CUG-BP1. *Oncogene* 35, 2087-2097
 18. Rong, Y., Liu, M., Liang, H., Guo, S., Xu, G., Min, Y., Lian, H., Xiang, Y., Zhang, S., and Xi, C. (2016) miR-138-5p contributes to cell proliferation and invasion by targeting Survivin in bladder cancer cells. *Molecular Cancer* 15, 82
 19. Althoff, K., Lindner, S., Odersky, A., Mestdagh, P., Beckers, A., Karczewski, S., Molenaar, J. J., Bohrer, A., Knauer, S., and Speleman, F. (2015) miR-542-3p exerts tumor suppressive functions in neuroblastoma by downregulating Survivin. *International journal of cancer* 136, 1308-1320
 20. Tochigi, H., Kajihara, T., Mizuno, Y., Mizuno, Y., Tamaru, S., Kamei, Y., Okazaki, Y., Brosens, J. J., and Ishihara, O. (2017) Loss of miR-542-3p enhances IGFBP-1 expression in decidualizing human endometrial stromal cells. *Scientific Reports* 7, 40001
 21. Zhao, S. (2013) Responsespecific progesterin resistance in a newly characterized Ishikawa human endometrial cancer subcell line resulting from longterm exposure to medroxyprogesterone acetate. *Oncology Letters* 5, 139-144
 22. Althoff, K., Lindner, S., Odersky, A., Mestdagh, P., Beckers, A., Karczewski, S., Molenaar, J. J., Bohrer, A., Knauer, S., and Speleman, F. (2015) miR-542-3p exerts tumor suppressive functions in neuroblastoma by downregulating Survivin. *International Journal of Cancer Journal International Du Cancer* 136, 1308
 23. Yoon, S., Choi, Y. C., Lee, S., Jeong, Y., Yoon, J., and Baek, K. (2010) Induction of growth arrest by miR-542-3p that targets survivin. *Febs Letters* 584, 4048-4052
 24. Karreth, F. A., and Pandolfi, P. P. (2013) ceRNA crosstalk in cancer: when ce-bling rivalries go awry. *Cancer Discovery* 3, 1113-1121
 25. Li, Q., Zhang, C., Chen, R., Xiong, H., Qiu, F., Liu, S., Zhang, M., Wang, F., Wang, Y., and Zhou, X. (2016) Disrupting MALAT1/miR-200c sponge decreases invasion and migration in endometrioid endometrial carcinoma. *Cancer Letters* 383, 28-40